



adonis laboratories pvt. ltd. — blending local resourcefulness with a global reach

Coniderm[®]

For Simple Fungal
Infections



CONIDERM Range

Description

Coniderm, **Coniderm-F** and **Coniderm-FG** are skin preparations targeted respectively to tackle early dermatomycoses (fungal infections), eczematous dermatomycoses and complicated dermatomycoses. Since all fungal infections have to be treated for a fairly long time, Coniderm Range is cosmetically acceptable as it has been formulated with a base that has the following properties: -

- soft and smooth texture
- easy spreadability
- non-staining
- lingering lilac fragrance

CONIDERM Topical Antifungal Cream

Composition

Miconazole Nitrate IP 2% w/w
in an easy to spread cream base.

Indications

Uncomplicated early Dermatophytoses (Tineas) caused by Trichophyton, Epidermophyton, Microsporum species as well as candida and pityrosporum.

Mode of Use

Patients should be advised to wash and dry the affected area, apply sufficient cream and gently rub it in. Coniderm should be used in the morning and evening.

Coniderm-F®

For Florid Dermatomycoses



Availability

Tubes of 15 gms

CONIDERM-F

Topical Antifungal, Anti-inflammatory Cream

Composition

Miconazole Nitrate IP	2% w/w
Fluocinolone Acetonide IP	0.01% w/w

in an easy to spread cream base.

Indications

Eczematous, itchy and inflamed lesions associated with all Dermatomycoses.

Mode of Use

Patients should be advised to wash and dry the affected area, apply sufficient cream and gently rub it in. Coniderm-F should be used in the morning and evening.

Availability

Tubes of 15 gms

Coniderm-FG®

For eczematous and infected fungal patches



CONIDERM-FG

Topical Antifungal, Antibacterial, Anti-inflammatory Cream

Composition

Miconazole Nitrate IP	2% w/w
Gentamicin Sulphate IP	
equivalent to Gentamicin base	0.1% w/w
Fluocinolone Acetonide IP	0.01% w/w

in an easy to spread cream base.

Indications

Chronic or complicated, inflamed and infected dermatoses of mixed aetiology.

Mode of Use

Patients should be advised to wash and dry the affected area. **Coniderm-FG** should be applied gently rubbed into the affected area twice or thrice daily for upto 3 to 6 weeks.

Availability

Tubes of 5 gms

All products are to be taken only under medical supervision.